

Numerical Simulations of the pulsed Jet of MWC 560

Matthias Stute & Max Camenzind

Landessternwarte Heidelberg, Königstuhl, D-69117 Heidelberg, Germany

Hans Martin Schmid

Institute of Astronomy, ETH Zentrum, CH-8092 Zurich, Switzerland

Abstract. MWC 560 (= V694 Mon) is the only known Symbiotic Star system in which the jet axis is practically parallel to the line of sight. Therefore this system is predestinated to study the dynamical evolution and the propagation of stellar jets. Spectroscopic monitoring done by Schmid et al. (2001) showed that the outflow is seen as absorption features in the continuum of the accretion disk and the accreting white dwarf, the emission line spectrum of the accretion disk and the spectrum of the red giant. We present the first numerical simulations of the jet of this particular object using the *NIRVANA* code (Ziegler & Yorke 1997) in order to reproduce the velocity structures seen in the observational data. This code solves the equations of hydrodynamics and was modified to calculate radiative losses due to non-equilibrium cooling by line-emission (Thiele 2000).

1. The Observations

The observational data revealed highly variable absorption structures which could be used to determine the velocity and acceleration structure of the jet. There exists a stable component with a radial velocity of $-1100 \pm 200 \text{ km s}^{-1}$ which represents the jet outflow at a larger distance. In addition, strong high velocity components with radial velocities up to -2500 km s^{-1} appear repeatedly on time scales of a few days and decay within several days. At the same time, the absorption between the absorption trough and the emission line with radial velocities of ≈ -900 to -400 km s^{-1} disappears which is an indication for an acceleration process near the jet source (see dynamical spectra in Schmid et al. 2001).

2. The Models

Due to the enormous computational costs, only 2D-models with the assumption of axisymmetry (2.5D) have been calculated yet. Therefore the companion became physically a Red Giant “Ring”. First, we started a purely hydrodynamical model without cooling. The parameters of this pulsed jet model are listed in Table 1. The influence of the stellar wind to the surrounding medium was

simulated with an $1/r^2$ law. The gravitational potential of the Red Giant and the Jet source was taken into account. The jet pulses were simulated by an enhancement of the jet velocity in the nozzle to 2000 km s^{-1} occurring every seventh day and lasting one day. Thereby also the jet density was varied to 0.25 or 2 times the normal density.

Table 1. Parameters of the models

	parameter	value
jet	number density	$5 \cdot 10^6 \text{ cm}^{-3}$
	temperature	10000 K
	velocity	10^3 km s^{-1}
jet pulse	number density (models I & II)	$1.25 \cdot 10^6 \text{ cm}^{-3}$
	number density (model III)	$1.0 \cdot 10^7 \text{ cm}^{-3}$
	velocity (models I - III)	$2 \cdot 10^3 \text{ km s}^{-1}$
stellar wind	$\dot{M}_{\text{red giant}}$	$1.0 \cdot 10^{-6} \text{ M}_{\odot} \text{ yr}^{-1}$
	velocity	10 km s^{-1}
system parameters	jet-radius	1 AU
	radius of the Red Giant	1 AU
	binary separation	4 AU
simulation parameters	integration domain	$50 \times 30 \text{ AU}$
	numerical resolution	20 grid cells/AU

In a second simulation, a cooling mechanism was implemented. According to Sutherland & Dopita (1993), three species (electrons, protons and hydrogen atoms) were considered and the heavier elements were simulated with a general cooling curve. In a third – again purely hydrodynamical – simulation, the number density of the jet pulse was increased to eight times the value of the first two models.

3. Results

While the complete hydrodynamical simulation (450 days) took two weeks on a workstation, after nine weeks on a NEC SX5 vector computer only 74 days of the simulation with cooling were calculated. These computational expenses are caused by the unique parameter choice, especially the high densities, resulting in a small cooling time step. These are the first simulations in this parameter region including cooling.

Comparing the logarithm of the total density for the same time step, one realizes the effects of cooling: due to smaller pressures in the jet the radial extension of the jet, the cross section and the resistance exerted by the external medium are lessened and therefore the cooled jet propagates faster (Fig. 1).

Using the density and velocity data, theoretical line profiles can be calculated, assuming the absorption $I = I_0 e^{-\tau}$ for each grid cell with

$$\tau_{\lambda} = \frac{\pi e^2}{m_e c} \lambda N_j f_{jk}. \quad (1)$$




Figure 1. Logarithm of density after 74 days; without (left) and with cooling (right)

With the increased number density of the jet pulse, the third simulation is able to reproduce the measured absorption of the fast pulse components (Fig. 2).

4. Discussion

In the purely hydrodynamical model the high velocity component decays much faster than in the observations – on time scales of one day. Taking into account the effects of cooling, it should decay slower which would be in agreement with the observational data. Due to the increasing propagation velocity of the jet and the decreasing resistance, the faster component should be able to exist longer. As this simulation is just at its beginning – and the jet has not yet reached its quasi-stationary phase, the results are still preliminary. But one first result could be the importance of cooling effects for modeling the jet correctly.

A parameter study varying further the density and the injection velocity during the outbursts should bring deeper insights into the real mass loss rate and jet energy in this system.

References

- Sutherland, R. S., & Dopita, M. A. 1993, *ApJS*, 88, 253
 Ziegler, U., & Yorke, H. 1997, *Comp. Phys. Comm.*, 101, 54
 Thiele, M. 2000, PhD Thesis, University of Heidelberg
 Schmid, H. M., Kaufer, A., Camenzind, M., Rivinius, Th., Stahl, O., Szeifert, T., Tubbessing, S., Wolf, B. 2001, *A&A*, 377, 206




Figure 2. Jet absorption line profiles for a series of eight consecutive days covering a high velocity pulse; theoretical (left) and observed (right)